


ENZYMATIC TEST KIT FOR THE DETERMINATION OF AMMONIA IN GRAPE JUICE AND WINE

PRODUCT

Product no. 4A120, for 30 tests, for *in vitro* use only

PRINCIPLE OF MEASUREMENT

Ammonia is found in both grape juice and wine. It is an important nutrient for yeast during primary fermentation. It is determined enzymatically according to the following equation:



Ammonia reacts with α -ketoglutarate and reduced nicotinamide adenosine dinucleotide (NADH) in the presence of glutamate dehydrogenase (GIDH) to form L-glutamate and NAD. The amount of NADH consumed is measured at 340 nm and is stoichiometrically related to the amount of ammonia present. The **Yeast Available Nitrogen (YAN)** content of the juice can be determined by adding the Ammonia Nitrogen (AN) content to the Primary Amino Acid Nitrogen (PAAN) content. PAAN can be determined by Vintessential Laboratories Analysis Kit 4A110.

CONTENTS

The kit includes the following reagents:

Reagent No.	Reagent	Preparation	Quantity	Stability
1	Buffer	Nil	33 mL	2 years at 4°C
2	NADH	Add 1.7 mL of distilled water to either bottle as required, mix to dissolve	2 x 1.7 mL	2 years at 4°C (1 month at 4°C once diluted)
3	GIDH	Mix gently by inversion before use	0.7 mL	2 years at 4°C
4	Standard	Nil	3.3 mL	2 years at 4°C

The shelf life of Reagent 1 can be extended by placing aliquots in a freezer.

Do not freeze reagents 2 or 3.

Failure to store reagents at the recommended temperature will reduce their shelf life.

For concentration of Standard, refer to label on bottle.

SAFETY

- **Wear safety glasses**
- **Do not ingest Buffer or Standard as they contain sodium azide as a stabilizer**

PROCEDURE

Operating Parameters

Wavelength

340 nm

Cuvettes

1cm, quartz, silica, methacrylate or polystyrene

Temperature

20 – 25°C

Final volume in cuvette

3.12 mL

Zero

against air without cuvette in light path

SAMPLE PREPARATION

Samples should be diluted with distilled water to ensure concentration in the assay solution is no more than 40 mg/L (ppm). Ideally, A_1 should lie between 0.90 – 1.20 absorbance units.

For samples with less than 200 mg/L of Ammonia, a 1 in 5 dilution should be sufficient.

For samples containing between 200 – 400 mg/L of Ammonia, a 1 in 10 dilution would be appropriate.

Undiluted red wines or highly coloured undiluted juice samples require decolourisation.

To decolourise, add approximately 0.1 g of PVPP to 5 mL of sample in a test tube. Shake well for about 1 minute. Clarification is achieved by settling or filtering through Whatman No. 1 filter paper.

SAMPLE ANALYSIS

a. Pipette the following volumes of reagents into the cuvettes:

Reagent	Blank	Standard	Sample
1. Buffer	1.00 mL (1000 μ L)	1.00 mL (1000 μ L)	1.00 mL (1000 μ L)
2. NADH	0.10 mL (100 μ L)	0.10 mL (100 μ L)	0.10 mL (100 μ L)
Distilled water	2.00 mL (2000 μ L)	1.90 mL (1900 μ L)	1.90 mL (1900 μ L)
Sample/Standard		0.10 mL (100 μ L)	0.10 mL (100 μ L)

b. Mix well and read absorbances, A_1 , once constant (approximately 5 minutes).

c. Pipette the following reagent into the cuvettes:

3. GIDH	0.02 mL (20 μ L)	0.02 mL (20 μ L)	0.02 mL (20 μ L)
---------	----------------------	----------------------	----------------------

d. Mix well and read absorbances, A_2 , once reaction is complete (approx 20 minutes).

CALCULATIONS*

1. Calculate the Net Absorbance for the Blank, Sample and Standard:

$$\text{Net Absorbance, } A_N = A_1 - A_2$$

2. Calculate the Corrected Absorbance by subtracting the Net Absorbance for the Blank from the Net Absorbance for the Sample.

$$\text{Sample Corrected Absorbance, } A_C = \text{Sample } A_N - \text{Blank } A_N$$

3. Do the same for the Standard by substituting the Standard absorbance values in place of the Sample absorbance values.

4. Calculate the Ammonia concentration as follows;

$$\text{Ammonia (mg/L)} = A_C \times 84.3 \times \text{Dilution Factor}$$

5. Calculate Ammonia Nitrogen as follows;

$$\text{Ammonia Nitrogen (mg/L)} = \text{Ammonia (mg/L)} \times 0.82$$

To calculate YAN (Yeast Assimilable Nitrogen), simply add Ammonia Nitrogen (AN) to the Primary Amino Acid Nitrogen (PAAN) calculated from kit 4A110:

$$\text{YAN} = \text{PAAN} + \text{AN}$$

*A calculation spreadsheet is available for download at:

<http://www.vintessential.com.au/certification/calculation-worksheets/>

REFERENCES

1. Bergmeyer, H.U. *et al* 1985, *Methods of Enzymatic Analysis*, 3rd ed., vol. 8, pp. 454-461; Verlag Chemie, Weinheim.

© Copyright 2018, Vintessential Laboratories. All rights reserved. No part of this publication may be copied or reproduced by any means without the written permission of Vintessential Laboratories.